
Australia
turnover
species
southwest
Europe
northern
distance
diversity
dispersal
GDM
geographical
environmental
beta
gradients
variables
precipitation
patterns
linear
important
ecological
along
compositional
flora
space
regions
models
variation
dissimilarity
plants
rate
explained
temperature
soil
greater
gradient
figure
rates
relationships
decay
partial
supplementary
expected
purely
unique
electronic
proportion
regression
maximum
importance
used
deviance
predictor
climate
taxa
considered
environment
variable
accommodate
predictors
vertebrate
explaining
rspb.royalsocietypublishing.org
see
region
climatic
composition
August
differences
winter
constant
also
floras
spatial
compare
means
relationship
vary
predicted
passive
less
common
two
data
relative
among
modes
fitted
obtained
long-distance
morphology
contribution
material
total
function
contrast
whereas
phosphorus
entire
observed
curvilinear
seed
approach
indicates
site
studies
versus
multiple
extent
changes
magnitude
found
significant
functions
sites
different
quantify
rapid
holding
wind
similarity
subregion
change
using
associated
datasets
height
table
shared
ii
determining
Soc
non-stationarity
effects
identified
ant
plant
adaptations
Downloaded
ability
l-splines
nonlinear
Proc
cells
modelling
mode
history
km
l-spline