found
similar
residuals
sub-fossil
similarity
column
part
variables
provide
may
negatively
species
depth
cladocerans
sphaericus
counted
macrophyte
Jackson
spp
change
assemblages
studies
number
taxa
identified
also
Frey
related
relationship
associated
sampling
deepest
Thus
abundance
sp
ordinations
agreement
correlation
important
longirostris
movement
ordination
past
significant
size
high
joint
PVI
fish
direction
rotation
cladoceran
study
salinity
assemblage
body
test
lakes
PCA
close
RDAs
Davidson
Procrustes
Mu
information
length
significance
ephippia
concordance
variation
two
chitinous
whereas
key
respectively
nets
remains
Furthermore
arrows
However
shallow
multiple
applied
lu
water
samples
supported
Bekliog
TP
good
lm
therefore
fossil
modern
variance
order
plant
Lake
sedimentary
Alona
Daphnia
represented
PROTEST
ller
Lakes
Table
scores
pelagic
analysis
littoral
surface
showed
richness
community
sample
Mediterranean
degree
response
relationships
arrow
sediments
data
sediment
higher
sampled
including
Jeppesen
accord
total
Cladocera
show
absent
analyses
low
used
diversity
datasets
using
Nevalainen
evenness
distribution
lake
environmental
variable
employed
RDA