

habitats

densities

predation

tundra

visibility

vegetation strategies

risk squirrels

boreal

ground

squirrel

Krebs

local

vigilance

behaviour

observed

Appendix

alpine

time

transitions

GUDD

foraging

forest

Wheeler

rate

excluded

considered

across

Behaviour

different

Canada

higher

Gillis

visibility

vegetation

strategies

three

boreal

cover

et

whether

juveniles

dense

Study

five

peanut

effects

determining

high-visibility

open

Appendix

species

structure

important

time

particularly

relationship

food

role

arch

role

Donker adults

Wheeler

populations

Falco

excluded

quitting

juvenile

via

lower

high

decline

Although

spent

analysis

patch

sex

Habitat

Animal

shrub

effect

arctic

Department

low-visibility

trial

associated

age

individual

Fig

shrub-dominated

shrub-tundra

costs

distance

expected

individuals

canopy-forming

Hik

measured

Table

change

higher

Gillis

high

decline

three

less

Boonstra

ground

Karels

achieve

encroachment

mass

two

increase

northern

June

al

total

elevations

Byrom

affected

Salix

Shrub

shrub-dominated

shrub-tundra

breadth

population

greater

appeared

Donker

Morrison

populations

responses

change

higher

Gillis

high

decline

three

less

Boonstra

ground

Karels

achieve

encroachment

mass

two

increase

northern

June

al

total

elevations

Byrom

affected

Salix

Shrub

shrub-dominated

shrub-tundra

breadth

population

greater

appeared

Donker

Morrison

populations

responses

change