fish
Lake
lakes
biomanipulation
phytoplankton
Jeppesen
effects
biomass
lake
restoration
may
manipulation
zooplankton
water
species
macrophytes
removal
nutrient
plants
submerged
warm
temperate
Sediment
Biomanipulation
macrophyte
Restoration
Denmark
studies
production
reduction
Eutrophication
loading
higher
Combat
Table
abundance
mussels
increased
Fig
clarity
TP
chemical
evenness
total
depth
years
mg
high
control
effect
shift
shallow
large
potential
carp
increase
cyanobacteria
methods
external
Tool
used
turbid
small
well
Erik
community
richness
measures
Secchi
including
success
Chl
genera
following
basins
biodiversity
growth
benthic
changes
grazing
long-term
Daphnia
due
mean
quality
plant
also
state
clear
However
experiments
observed
cascade
Engelsholm
Vang
three
found
size
strong
therefore
low
reduced
trophic
Figure
alternative
decreased
lower
key
via
combined
abundant
perch
change
less
Fish
tilapia
biological
pelagic
composition
needed
L1
case
thus
TN
Sondergaard
area
two
eutrophic
several
Faarup
stocking
predation
pressure
reduce
summer
recovery
areas
increasing
piscivorous
cladocerans
number
Meerhoff
zebra
piscivores
typically
often
turbid
g.a
kg