

argan
woodlands
climate
distribution
range
woodland
Africa
climatic
areas
change
ecosystem
Mediterranean
Quaternary
human
Vegetation
current
North
Sahara
water
potential
precipitation
Science
spinosa
Morocco
dry
changes
southern
Petit
LGM
temperature
estimated
past
future
Desert
Benabid
Nieto-Lugilde
million
Maximum
region
northern
sensitivity
century
well
forests
also
Glacial
Atlantic
deforestation
semi-arid
tree
Fig
arid
Applied
Last
key
strongly
Le
high
response
Svenning
Argania
northwest
important
Benzyane
near
pressure
see
Holocene
particularly
strong
factors
losses
throughout
Mousadik
constitute
warming
tendency
Tarrier
conservation
findings
However
suitable
forest
10.1111/avsc.12133
contraction
studies
suggest
anchez
limit
Indeed
Morocco's
low
shift
ha
BP
severe
mid-Holocene
shifts
importance
indicate
order
north
year
Table
biodiversity
local
models
ecosystems
scenarios
results
likely
Alba-S
relatively
impact
provide
stress
latitudes
vulnerability
extended
species
Sapotaceae
coast
African
role
El
e.g
Western
International
mediterranean
aez
assess
Atlas
area
late
palaeorecords
SDM
Association
evidence
may
livelihoods
result
degradation
larger
already
quarter
within