species
vegetation
Europe
Fig
forest
fossil
Western
MIS
climate
Northwestern
stages
shifts
range
bird
changes
found
distribution
temperature
conditions
site
cold
birds
tree
period
fossils
Palearctic
also
Table
well
study
warm
indicating
glacial
periods
dated
sites
years
studies
past
BP
habitat
Pleistocene
Lagopus
Eemian
Figure
Central
forests
woodland
relatively
Eastern
temperate
Turdus
Furthermore
responses
Weichselian
Pyrrhocorax
category
stage
mammals
distributions
time
climatic
trees
north
high
S1
degree
different
results
much
environments
south
proportions
Eurasian
Poland
present
occurrences
provided
European
presence
Betula
one
using
two
effects
record
analysis
warmer
open
Late
types
requirements
due
southern
categories
openness
similar
areas
full-glacial
probably
avian
like
observed
low
Late-Glacial
perdix
coniferous
Perdix
close
Southern
Tetrao
predominated
Owl
lagopus
boreal
interglacial
evidence
assemblages
response
analyses
compared
clear
even
LGM
Fossils
deciduous
subregion
overall
Alps
landscape
gradient
many
within
change
muta
proportion
stratigraphic
refugia
ranges
findings
identified
layers
e.g
general
excluded
reported
Last
database
sp
Lagopus
Central
Figure
Eemian
relative
Eastern
temperate
Turdus
Furthermore
responses
Weichselian
Pyrrhocorax
category
stage
mammals
distributions
time
climatic
trees
north
high
S1
degree
different
results
much
environments
south
proportions
Eurasian
Poland
present
occurrences
provided
European
presence
Betula
one
using
two
effects
record
analysis
warmer
open
Late
types
requirements
due
southern
categories
openness
similar
areas
full-glacial
probably
avian
like
observed
low
Late-Glacial
perdix
coniferous
Perdix
close
Southern
Tetrao
predominated
Owl
lagopus
boreal
interglacial
evidence
assemblages
response
analyses
compared
clear
even
LGM
Fossils
deciduous
subregion
overall
Alps
landscape
gradient
many
within
change
muta
proportion
stratigraphic
refugia
ranges
findings
identified
layers