
EXCLUDED
SITE
ECOSYSTEM
MEGAFAUNAL
LOWER
IMPORTANT
WITHIN
PROPORTION
CONSERVATION
COMPARE
SENSITIVITY
ANALYZED
PRESENT
TEMPORAL
SUPPORT
HUMANS
BEETLE
HABITATS
PNAS
DENSITY
GLACIAL
REFLECT
HIGH
RADIOMETRIC
WOOD
CONTRAST
S2
MINIMUM
LANDSCAPE
LESS
SMITH
BUGSCEP
WELL
DATABASE
REPRESENTATION
DOMESTIC
AGRICULTURAL
RANGE
BIOME
THERMAL
INTERGLACIAL
CLOSED
ONE
OPEN
GIVEN
MODERN
BEETLES
CATEGORY
STRUCTURE
LATE
PERIOD
ASSEMBLAGES
WEIGHT
BASED
WHITEHOUSE
TEMPERATURES
PLEISTOCENE
MANAGEMENT
RELATIVE
NAMES
2A
LOSSES
FOUR
HERBIVORES
SPECIES
SAMPLES
VEGETATION
FOUND
TEST
USED
MAXIMUM
ROLE
DENSITIES
ALSO
SIMILAR
HERBIVORE
RECORDED
LOW
CLIMATE
CATEGORIES
ANALYSIS
GREAT
SEMICLOSED
WOODLAND
ABUNDANCE
LARGE
HOLOCENE
TIME
USING
SAMPLES
INTERGLACIAL
SITES
KG
LAST
DUNG
TWO
GLMM
TREES
PRIMARILY
WILD
EARLY
MAY
S1
C.J.S
MEDIAN
FIRE
ASSIGNED
YEARS
ECOSYSTEMS
ASSOCIATED
DUNG
LANDSCAPES
LAST
ECOLOGICAL
DATA
RESULTS
HIGHER
VALUES
IMPORTANCE
FIG
PERIOD
FOREST
PERIODS
GREATER
COVER
B.P
INDICATED
PASTURE
DIVERSITY
DATES
MEGAFAUNA
RANDOM
DIVERSE
REPRESENT
EUROPE
INCLUDED
TEMPERATE
EARLY
INDICATE
SAMPLE
ABUNDANT
BRITAIN
SELECTED
EXTINCT
LATE
THRESHOLD
CALCULATED
CONDITIONS
HUMAN
PER